

# Burkholderia Cepacia Selective Agar, Complete

## SKU: 700003514, 700003515, 700003516

## NCM0209

#### Intended Use

Burkholderia Cepacia Selective Agar, Complete is used for the selective isolation of the *Burkholderia* (*Pseudomonas*) cepacia Complex (Bcc). Burkholdaria Cepacia Selective Agar is not intended for use in the diagnosis of disease or other conditions in humans

#### **Description**

The United States Pharmacopeia has updated the USP/NF Microbiological Test <60> Microbiological Examination of Nonsterile Products- Tests for *Burkholderia cepacia* Complex (Bcc) to include the use of this medium. This test is designed to determine whether a substance or preparation complies with an established specification for microbiological quality and/or to evaluate whether product contains members of the Bcc.

Burkholderia Cepacia Selective Agar, Complete is the complete medium formulation specified by the USP/NF. BCSA, Complete contains peptones and sugars that supply nutrients for the growth of the Bcc. Phenol Red is the sugar utilization indicator. Crystal Violet is added to inhibit the growth of Gram-positive bacteria. Antimicrobics are added to suppress or inhibit the recovery bacteria other than the Bcc. Agar is the solidifying agent.

#### Formula / Liter

| Enzymatic Digest of Casein | 10.0 g/L     |
|----------------------------|--------------|
| Lactose                    | 10.0 g/L     |
| Sucrose                    | 10.0 g/L     |
| Sodium Chloride            | 5.0 g/L      |
| Yeast Extract              | 1.5 g/L      |
| Phenol Red                 | 0.08 g/L     |
| Crystal Violet             | 0.002 g/L    |
| Vancomycin                 | 0.0025 g/L   |
| Gentamicin                 | 0.01 g/L     |
| Polymyxin B                | 600,000 IU/L |
| Agar                       | 14.0 g/L     |
| Final pH 6.8 ± 0.3 at 25°C |              |

Formula is adjusted or supplemented as required to meet performance specifications

#### **Precaution**

Refer to SDS

#### **Preparation**

- 1. Suspend 50.69 grams of the medium in one liter of purified water.
- 2. Heat with frequent agitation and bring to a boil. Avoid overheating.
- 3. Autoclave at 121°C for 15 minutes.
- 4. Cool to 45-50°C.
- 5. Avoid holding for longer than the time required to equilibrate.



620 Lesher Place • Lansing, MI 48912 800-234-5333 (USA/Canada) • 517-372-9200 foodsafety@neogen.com • foodsafety.neogen.com



## Test Procedure

Follow the USP Method <60> Microbiological Examination of Nonsterile Products- Test for *Burkholderia cepacia* Complex<sup>2</sup>.

### **Quality Control Specifications**

**Dehydrated appearance:** Powder is homogenous, free-flowing and light beige and may have a slight pink or gray tint.

Prepared appearance: The prepared medium is clear to slightly hazy and red-violet without a precipitate.

#### **Expected Cultural Response**

Cultural response on Burkholderia Cepacia Selective Agar, Complete when tested following the USP/NF <60> Surface Spread Method (Microbial Enumeration Tests <61>) Growth Promotion Test; Suitability of the Counting Method and Controls, Suitability of the Counting Method in the Presence of Product, Recovery of Microorganisms in the Presence of Product.

| MICROORGANISM  | APPROX.<br>INOCULUM<br>(CFU) | EXPECTED RESULTS        |                                |
|--|------------------------------|-------------------------|--------------------------------|
|  |                              | Growth                  | Reaction                       |
| <i>Burkholderia cepacia</i><br>ATCC <sup>®</sup> 25416 | 10-100                       | Growth                  | Yellow-Orange to Pink-Red Zone |
| <i>Burkholderia cepacia</i><br>ATCC <sup>®</sup> 25608 | 10-100                       | Growth                  | Yellow-Orange to Pink-Red Zone |
| Burkholderia cenocepacia<br>ATCC® BAA-245              | 10-100                       | Growth                  | Yellow-Orange to Pink-Red Zone |
| Burkholderia multivorans<br>ATCC <sup>®</sup> BAA-247  | 10-100                       | Growth                  | Yellow-Orange to Pink-Red Zone |
| Pseudomonas aeruginosa<br>ATCC® 9027                   | 10 <sup>3</sup>              | Suppressed to Inhibited |                                |
| Staphylococcus aureus<br>ATCC® 6538                    | 10 <sup>3</sup>              | Suppressed to Inhibited |                                |
| <i>Escherichia coli</i><br>ATCC <sup>®</sup> 25922     | 10 <sup>3</sup>              | Suppressed to Inhibited |                                |
| Enterococcus faecalis<br>ATCC <sup>®</sup> 29212       | 10 <sup>3</sup>              | Suppressed to Inhibited |                                |

The organisms listed are the minimum that should be used for quality control testing

### <u>Results</u>

Refer to appropriate references for test results.



620 Lesher Place • Lansing, MI 48912 800-234-5333 (USA/Canada) • 517-372-9200 foodsafety@neogen.com • foodsafety.neogen.com



## **Expiration**

Refer to the expiration date stamped on the container label. The dehydrated medium should be discarded if it is not free-flowing, or if the medium has changed from the original color. Expiry applies to the medium in its intact container when stored as directed.

## **Limitations of the Procedure**

Due to nutritional variations, some strains may be encountered that grow poorly or fail to grow on this medium.

## Storage

Store the dehydrated culture media at 2-8°C away from direct light. Once opened and recapped, place the container back in the same storage conditions. Protect from moisture and light by keeping container tightly closed.

## **References**

- 1. Henry, D. et al. 1997, Identification of *Burkholderia cepacia* Isolates from Patients with Cystic Fibrosis and Use of a Simple New Selective Medium. *J. Clin. Micro.:35:614-619.*
- 2. USP/NF <60> Microbiological Examination of Nonsterile Products- Test for *Burkholderia cepacia* Complex.



