

## Enterobacteria Enrichment (EE) Broth Mossel

**SKU: 700003125, 700003126, 700003127, 700003128, 700004438  
NCM0057**

### Intended Use

Enterobacteria Enrichment (EE) Broth Mossel is used for the cultivation and enrichment of *Enterobacteriaceae* in food. Enterobacteria Enrichment (EE) Broth Mossel, conforms to Harmonized USP/EP/JP Requirements and is not intended for use in the diagnosis of disease or other conditions in humans.

### Description

Enterobacteria Enrichment Broth Mossel was developed by Mossel, Visser, and Cornelissen to facilitate the growth of *Enterobacteriaceae*. This medium contains dextrose to enhance the growth of *E. coli* and *Salmonella* spp., particularly in food samples. Nuisance organisms are suppressed by the addition of Ox Bile and Brilliant Green.

Enterobacteria Enrichment Broth Mossel is used as an enrichment broth, providing a rich environment for the recovery of damaged or injured cells. *Enterobacteriaceae* organisms can be injured in food-processing procedures, including exposure to low temperature, sub-marginal heat, drying, radiation, preservatives, or sanitizers. The enumeration of *Enterobacteriaceae* is an important measure of the sanitary condition of food.

EE Broth, Mossel complies with the specifications of the Eiprodukte-Verordnung (German Egg Product Regulations) and conforms to Harmonized United States Pharmacopoeia (USP), European Pharmacopoeia (EU), and Japanese Pharmacopoeia (JP).

### Typical Formulation

Bile Salts	20.0 g/L
Enzymatic Digest of Gelatin	10.0 g/L
Disodium Hydrogen Phosphate Dihydrate	8.0 g/L
Glucose Monohydrate	5.0 g/L
Potassium DiHydrogen Phosphate,	2.0 g/L
Brilliant Green	0.015 g/L

pH: 7.2 ± 0.2 at 25°C

Formula is adjusted and/or supplemented as required to meet performance specifications.

### Precaution

Refer to SDS

### Preparation

1. Suspend 45g of the medium in one liter of purified water.
2. Heat at 100°C for 30 minutes to completely dissolve the medium.
3. Cool rapidly in cold water.
4. DO NOT AUTOCLAVE.

### Test Procedure

1. Inoculate prepared Enterobacteria Enrichment (EE) Broth Mossel with approximately 10 g of homogenized food or other material to be tested.
2. Shake the inoculated medium thoroughly for a few seconds to mix well.
3. Incubate for a total of 18 - 48 hours at 30 - 35°C. Shake tubes or flasks after the first 3 hours of incubation.



620 Leshar Place • Lansing, MI 48912  
800-234-5333 (USA/Canada) • 517-372-9200  
foodsafety@neogen.com • foodsafety.neogen.com

# Technical Specification Sheet



4. Streak a loopful of the incubated enrichment culture of Enterobacteria Enrichment (EE) Broth - Mossel onto a prepared selective medium.
5. Incubate the plates for 18 - 24 hours at 30 - 35°C. Examine the incubated medium for the presence of the target organism.

## **Quality Control Specifications**

**Dehydrated Appearance:** Powder is homogeneous, free-flowing, and light beige to light green.

**Prepared Appearance:** Prepared medium is clear to slightly hazy with no to trace precipitate and green to dark green in color.

**Expected Cultural Response and USP/EP/JP Growth Promotion:** Cultural response in Enterobacteria Enrichment Broth - Mossel at 30 - 35°C after 18 – 48 hours of incubation and subcultured onto Violet Red Bile Glucose Agar after 18-24 hours and 40-48hours. VRBGA plates were incubated at 30-35°C for 18-24 hours.

Microorganism	Approx. Inoculum (CFU)	Expected Results Growth
<i>Escherichia coli</i> ATCC® 8739	10-100	Recovered at >10 cfu
<i>Escherichia coli</i> ATCC® 25922	10-100	Recovered at >10 cfu
<i>Enterococcus faecalis</i> ATCC® 29212	~10,000	Complete inhibition
<i>Enterococcus faecalis</i> ATCC® 19433	~10,000	Complete inhibition
<i>Pseudomonas aeruginosa</i> ATCC® 9027	10-100	Recovered at >10 cfu
<i>Salmonella typhimurium</i> ATCC® 14028	10-100	Recovered at >10 cfu
<i>Staphylococcus aureus</i> ATCC® 6538	~10,000	Inhibited
<i>E.coli</i> ATCC® 8739 + <i>E. faecalis</i> ATCC® 19433	10-100 ~10,000	Recovered at >10 cfu Complete Inhibition
<i>E.coli</i> ATCC® 8739 + <i>E. faecalis</i> ATCC® 29212	10-100 ~10,000	Recovered at >10 cfu Complete Inhibition

The organism listed are the minimum that should be used for quality control testing.

## **Results**

Examine Enterobacteria Enrichment (EE) Broth Mossel for growth, indicated by turbidity.

## **Expiration**

Refer to expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color.

## **Limitation of the Procedure**

Some strains may be encountered that grow poorly or fail to grow on this medium.

## **Storage**

Store dehydrated culture media at 2 – 30°C away from direct sunlight. Once opened and recapped, place the container in a low humidity environment at the same storage temperature. Protect from moisture and light by keeping container tightly closed.



620 Leshar Place • Lansing, MI 48912  
800-234-5333 (USA/Canada) • 517-372-9200  
foodsafety@neogen.com • foodsafety.neogen.com

## References

1. European Pharmacopoeia 10<sup>th</sup> Edition (2020)
2. United States Pharmacopeia National Formulary 2018: USP 41 NF 36
3. Japanese Pharmacopeia 17<sup>th</sup> Edition (2017)
4. Mossel, Vissar, and Cornellisen. 1963. J. Appl. Bacteriol. 26:444.
5. Hartman, P. A., and S. A. Minnich. 1981. Automation for rapid identification of salmonellae in foods. J. Food Prot. 44:385-386.
6. Sorrells, K. M., M. L. Speck, and J. A. Warren. 1970. Pathogenicity of *Salmonella gallinarum* after metabolic injury by freezing. Appl. Microbiol. 19:39-43.
7. Bundesminister für Jugend, Familie und Gesundheit: 1975. Verordnung über die gesundheitlichen Anforderungen an Eiprodukte und deren Kennzeichnung (Eiprodukte-Verordnung).

Record ID: FS-TSS-0231 Revision Number: 3.0 Effective Date: 2023-10-24 12:00 AM EDT