

**Middlebrook 7H11 Agar**  
**SKU: 700003073, 700003074, 700003075, 700003076**  
**NCM0043**

## **Intended Use**

Middlebrook 7H11 Agar is used with glycerol and OADC Enrichment for the cultivation of *Mycobacterium* spp. in a laboratory setting. Middlebrook 7H11 Agar is not intended for use in the diagnosis of disease or other conditions in humans.

## **Description**

There are two types of solid culture media for the primary isolation of mycobacteria, coagulated egg as a base (Lowenstein formulations) and an agar base (Middlebrook formulations). The use of agar-based media for primary isolation of mycobacteria have the following significant advantages:

1. Agar-based media do not usually liquefy in the presence of contaminating proteolytic organisms.
2. Agar-based media retain exact concentrations of added drugs because the medium is solidified with agar rather than by inspissation of the egg. There is less drug inactivation when egg ingredients are absent.

Middlebrook 7H11 Agar is a modification of Middlebrook 7H10 Agar Special as recommended by Cohn, Waggoner, and McClately. Cohn et al. added an enzymatic digest of casein and found organism growth was stimulated for fastidious strains of *Mycobacterium tuberculosis*.

## **Typical Formulation**

|                            |             |
|----------------------------|-------------|
| Enzymatic Digest of Casein | 1.0 g/L     |
| Disodium Phosphate         | 1.5 g/L     |
| Monopotassium Phosphate    | 1.5 g/L     |
| Ammonium Sulfate           | 0.5 g/L     |
| Monosodium Glutamate       | 0.5 g/L     |
| Sodium Citrate             | 0.4 g/L     |
| Ferric Ammonium Citrate    | 0.04 g/L    |
| Magnesium Sulfate          | 0.05 g/L    |
| Copper Sulfate             | 0.001 g/L   |
| Pyridoxine                 | 0.001 g/L   |
| Zinc Sulfate               | 0.001 g/L   |
| Biotin                     | 0.0005 g/L  |
| Malachite Green            | 0.00025 g/L |
| Agar                       | 13.5 g/L    |

## **Supplement**

Glycerol, 5 mL  
OADC Enrichment, 100 mL

Final pH: 6.6 ± 0.2 at 25°C

Formula is adjusted and/or supplemented as required to meet performance specifications.

## **Precaution**

Refer to SDS

## **Preparation**

1. Suspend 19 g of the medium in 900 mL of purified water containing 5 mL of glycerol.
2. Heat to boiling to dissolve completely.
3. Autoclave at 121°C for 10 minutes.
4. Cool to 45 - 50°C and aseptically add 100 mL of OADC Enrichment.

## **Quality Control Specifications**

**Dehydrated Appearance:** Powder is homogeneous, free flowing, and beige.

# Technical Specification Sheet



**Prepared Appearance:** Prepared medium is trace to slightly hazy and pale to light green to gray-white.

**Expected Cultural Response:** Cultural response on Middlebrook 7H11 Agar at incubated under CO<sub>2</sub> at 35 ± 2°C and examined for growth after 3 – 28 days incubation.

| Microorganism   | Approx. Inoculum (CFU) | Expected Results |
|---|------------------------|------------------|
| <i>Mycobacterium fortuitum</i> Group IV ATCC® 6841        | Heavy                  | Growth           |
| <i>Mycobacterium intracellulare</i> Group III ATCC® 13950 | Heavy                  | Growth           |
| <i>Mycobacterium kansasii</i> Group I ATCC® 12478         | Heavy                  | Growth           |
| <i>Mycobacterium scrofulaceum</i> Group II ATCC® 19981    | Heavy                  | Growth           |
| <i>Mycobacterium tuberculosis</i> H37Ra ATCC® 25177       | Heavy                  | Growth           |

The organisms listed are the minimum that should be used for quality control testing.

## **Test Procedure**

Inoculate specimen onto the medium. Incubate tubes for up to eight weeks. Examine tubes for growth at regular intervals. Refer to specific procedures for a complete discussion on the isolation and identification of *Mycobacterium* spp.

## **Results**

Observe colonies that may or may not be pigmented. Colony morphology is dependent on the species isolated.

## **Expiration**

Refer to expiration date stamped on the container. Dehydrated medium should be discarded if not free flowing, or if appearance has changed from original color. Expiry applies to medium in its intact container when stored as directed.

## **Limitations of the Procedure**

Due to nutritional variation, some strains may be encountered that grow poorly or fail to grow on this medium. Further tests are necessary for confirmation of *Mycobacterium* spp.

## **Storage**

Store sealed bottle containing dehydrated medium at 2 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light by keeping container tightly closed.

## **References**

1. Musser, J. M. 1995. Antimicrobial resistance in Mycobacteria: molecular genetic insights. Clinical Microbiology Reviews. 8:496-514.
2. Murray, P. R., E. J. Baron, M. A. Pfaller, F. C. Tenover, and R. H. Tenover (eds.). 1995. Manual of clinical microbiology, 6th ed. American Society for Microbiology, Washington, D.C.
3. Isenberg, H. D. (ed.). 1992. Clinical microbiology procedures handbook, vol. 1 American Society for Microbiology, Washington, D.C.
4. Cohn, M. L., R. F. Waggoner, and J. K. McClatchy. 1968. The 7H11 Medium for the cultivation of mycobacteria. Am. Rev. Resp. Dis. 98:295.

