

Technical Specification Sheet



Alkaline Saline Peptone Water (ASPW) (ISO) (NCM0175)

Intended Use

Alkaline Saline Peptone Water is a medium for the enrichment of *Vibrio* spp. from food and water samples according to ISO 21872-1:2017, and is not intended for use in the diagnosis of disease or other conditions in humans.

Description

Originally described by Shread, Donovan & Lee as an enrichment broth for *Aeromonas* spp. and identified by Cruickshank as an effective medium for the enrichment of *Vibrio* spp., Alkaline Saline Peptone Water uses elevated pH and salt levels to provide a favorable environment for the enrichment of *Vibrio* spp.

Peptone provides essential vitamins, minerals, amino acids & nitrogen for growth requirements. Sodium chloride provides essential electrolytes for maintenance of the osmotic balance.

Sub-culture onto Thiosulfate Citrate Bile Salts Sucrose (TCBS) Agar (NCM0052) will be required.

Typical Formulation

Peptone	20.0 g/L
Sodium Chloride	20.0 g/L

Final pH: 8.6 ± 0.2 at 25°C

Formula may be adjusted and/or supplemented as required to meet performance specifications.

Precaution

Refer to SDS

Preparation

1. Dissolve 40 grams of the medium in one liter of purified water.
2. Heat with frequent agitation to completely dissolve the medium, if necessary.
3. Autoclave at 121°C for 15 minutes.

Test Procedure

For a complete discussion on the isolation and identification of *Vibrio cholerae* and other enteropathogenic *Vibrio* spp., refer to specific procedures, such as ISO 21872-1:2017.

Quality Control Specifications

Dehydrated Appearance: Powder is homogeneous, free flowing and beige.

Prepared Appearance: Prepared medium is a clear, very pale-yellow liquid.

Expected Cultural Response: Cultural response in ASPW at 36-38°C after 16-20 hours of incubation.

Microorganism	Approx. Inoculum (CFU)	Response
<i>Vibrio cholera</i> ATCC® 14733	10-100	Growth
<i>Vibrio furnissii</i> NCTC 11218	10-100	Growth
<i>Vibrio parahaemolyticus</i> NCTC 10885	10-100	Growth



620 Lesher Place • Lansing, MI 48912
 800-234-5333 (USA/Canada) • 517-372-9200
 foodsafety@neogen.com • foodsafety.neogen.com

Technical Specification Sheet



Results

After sub-culture of the ASPW onto TCBS Agar, with 18 – 48 hours of incubation at 36-38°C, sucrose-fermentating *Vibrio* spp. (*V. cholerae*, *V. alginolyticus*, *V. hareyi*, *V. cincinnatiensis*, *V. fluvialis*, *V. furnissii*, *V. metschnikovii*, and some *V. vulnificus*) appear smooth, opaque, thin-edged yellow colonies on TCBS Agar.

Expiration

Refer to expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing or appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedures

Due to nutritional variation, some strains may be encountered that grow poorly or fail to grow on this medium.

Storage

Store dehydrated culture media at 2-30°C away from direct sunlight. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light by keeping container tightly closed.

References

1. Cruickshank R. (1968) Medical Microbiology. 11th ed. Livingstone Ltd, London, UK.
2. ISO 21872-1:2017 Microbiology of the food chain – Horizontal method for the determination of *Vibrio* spp. – Part 1: Detection of potentially enteropathogenic *Vibrio parahaemolyticus*, *Vibrio cholerae* and *Vibrio vulnificus*.
3. Shread, P., Donovan, T.J. and Lee, J.V. (1991). *Soc. Gen. Microbiol. Q.* 8. 184.



620 Leshner Place • Lansing, MI 48912
800-234-5333 (USA/Canada) • 517-372-9200
foodsafety@neogen.com • foodsafety.neogen.com