

For Laboratory Use

Store at 2–8°C



E. Coli Supplement (MUG) Part #S2-ECM

For use with S2-EC only
12 mL per vial

Usage Instructions:

1. Swirl supplement prior to use. If crystals are observed, warm gently until dissolved.
2. Add 0.120 mL of the S2-ECM to the *E. coli* (S2-EC) test vial.
3. Place S2-ECM back in the refrigerator.



620 Leshler Place, Lansing, MI 48912
800/234-5333 (USA/Canada) or 517/372-9200 • fax: 517/372-2006
e-mail: foodsafety@neogen.com • www.neogen.com

© Neogen Corporation, 2018. Neogen and Soleris are registered trademarks of Neogen Corporation.
C0120126A

S2-ECM-0318

For Laboratory Use

Store at 2–8°C



E. Coli Supplement (MUG) Part #S2-ECM

For use with S2-EC only
12 mL per vial

Usage Instructions:

1. Swirl supplement prior to use. If crystals are observed, warm gently until dissolved.
2. Add 0.120 mL of the S2-ECM to the *E. coli* (S2-EC) test vial.
3. Place S2-ECM back in the refrigerator.



620 Leshler Place, Lansing, MI 48912
800/234-5333 (USA/Canada) or 517/372-9200 • fax: 517/372-2006
e-mail: foodsafety@neogen.com • www.neogen.com

© Neogen Corporation, 2018. Neogen and Soleris are registered trademarks of Neogen Corporation.
C0120126A

S2-ECM-0318

For Laboratory Use

Store at 2–8°C



E. Coli Supplement (MUG) Part #S2-ECM

For use with S2-EC only
12 mL per vial

Usage Instructions:

1. Swirl supplement prior to use. If crystals are observed, warm gently until dissolved.
2. Add 0.120 mL of the S2-ECM to the *E. coli* (S2-EC) test vial.
3. Place S2-ECM back in the refrigerator.



620 Leshler Place, Lansing, MI 48912
800/234-5333 (USA/Canada) or 517/372-9200 • fax: 517/372-2006
e-mail: foodsafety@neogen.com • www.neogen.com

© Neogen Corporation, 2018. Neogen and Soleris are registered trademarks of Neogen Corporation.
C0120126A

S2-ECM-0318

For Laboratory Use

Store at 2–8°C



E. Coli Supplement (MUG) Part #S2-ECM

For use with S2-EC only
12 mL per vial

Usage Instructions:

1. Swirl supplement prior to use. If crystals are observed, warm gently until dissolved.
2. Add 0.120 mL of the S2-ECM to the *E. coli* (S2-EC) test vial.
3. Place S2-ECM back in the refrigerator.



620 Leshler Place, Lansing, MI 48912
800/234-5333 (USA/Canada) or 517/372-9200 • fax: 517/372-2006
e-mail: foodsafety@neogen.com • www.neogen.com

© Neogen Corporation, 2018. Neogen and Soleris are registered trademarks of Neogen Corporation.
C0120126A

S2-ECM-0318